KOLKATA WEDNESDAY 29 MAY 2013 The Statesman

Cocking an eye for danger

The lowly rat makes good use of its two eves. savs s ananthanaravanan

NATURE has provided living things with pairs of eyes for good reason. A single eye can provide sight, but for accurate estimation of distance and depth we need two eyes that are placed some

two eyes that are placed some distance apart. But not all animals use both eyes all the time, and many use their two eyes singly, to command a wider field of view. Damian J Wallace, David S Greenberg, Juergen Sawinski, Stefanie Rulla, Giuseppe Notaro and Jason ND Kerr, at the Max Planck Institute for Biological Chyenretics and Biological Cybernetics and the Bernstein Center for Computational Neuroscience in Germany, report in the

journal Nature that rodents effectively combine both kinds of use of eyes to its best advantage.

The animal eye is nothing but a biological optical camera. It focuses light that

camera. It focuses light that falls upon it onto the retina, which is a light-sensitive screen. In other words, the wordd in front of the camera, which consists of objects at different distances from the eye, is captured on a flat surface, which has length and breadth but no depth. Large objects at a distance then look the same size as a look the same size as a smaller object that is closer, and a single eye, or any number of eyes, by

themselves cannot make out which object is nearer or further away. This is quite noticeable in photographs of landscapes or other scenes with depth — the awareness of distances gets obscured in

of distances gets obscured in the picture.

But with a pair of eyes, each sees a different picture, each from a different point, a few inches to the side. The brain is then able to merge these two pictures and bring with a presention of out the perception of distance. The views of the bicycle, house and tree, seen by each eye, one a little to the side, are shown in the second

If the reader presses one eye, so that the eyes get misaligned and there are two images, and four houses, he/she could try to merge the two in the middle — it will spring to life, with the tree clearly behind the house.



Another experiment to try is

going too far or too close perception. Such perception of depth, which is called *stereoscopic* vision, is what we get thanks to our two eyes. All animals that have two eyes are able to

Seen by right eye

correctly get to food or prey, avoid obstacles and cross barriers, thanks to this perception. But for the eyes to act in this way, there is also the need for both eyes to coordinate and align themselves to gaze at the same object at the same mobject at the same time, even when we move our head or shift our direction of gaze or change the focus to a different distance. This takes muscular coordination and is muscular coordination and is learnt in a few weeks by human babies, for instance, when they experiment with grasping or touching.

Using eyes singly Coordinated use of eyes for stereoscopic vision is the rule with larger animals and with larger animals and certainly with primates, like the chimpanze or humans. Keeping the eyes working together is useful for activities like leaping, hunting or using tools. But it involves a price, in limiting the field of attention. In the case of grazing birds, like the chicken, it is more useful if the bird could view all sides. the bird could view all sides at once to be able to locate things to eat. The chicken has, hence, evolved to have

whole yard at once.

Once a tidbit is located, the bird could sight the morsel with both eyes by turning its head this way and that, to capture two images which the brain could process and help the chicken to peck accurately.

The authors of the paper in Vetture note; that although

Nature note that although rodents share with other mammals the key mammals the key components of eye movement control, it is not known that rodents coordinate the movements of the eye even when they have not focused on an object or when they are moving freely. The group of researchers carried out a series of experiments where movement of the eyeball of freely moving rats was monitored with the help of head-mounted cameras and the position and orientation of the head itself was observed with the help of a specially designed tracking specially designed tracking

specially designed tracking system.

The movements of the heatest wave the roll, or tilting leaf and right, the pitch, or looking up and down, and the yaw, which is to turn to the left or right. While the head was moving, it was found that the eyes carried out wide excursions and were markedly not coordinated. Not only did the pupils move inward, outward, up and down, they even rotated about the direction of vision. And it was found that the And it was found that the

adout in execution to inson. And it was found that the movements of the pupils were dictated by the orientation of the head. Earlier studies have shown that rolling leads to raising of the cychall on the side that goes down, or vice-versa, and pitching leads to the cychalls moving to the centre for pitch up and outwards for the pitch down. And pitching motion also leads to rotation of the pupil. The movement of the pupils of free nowing rats was seen to mostly follow this rule.

free-moving rats are often not pointed in the same direction. This led to the unction. Inis ied to the matching images of the two eyes only in part of the retina and at different parts every instant. This was not as much the case when the head was clamped or restrained, or when the rat held its head still as when; it had to come a still, as when it had to cross a gap in the floor, but there was little evidence of coordination to keep a fixed relationship between the


cyes.

The result is that the rat has a large field of view, about 200°, in the front, above and to the rear, but most of the vision with the use of one eye only. The parts of the field where the images of the two eyes overlap are directly in front and above the head. Studies of the overlap for different orientations of the head show that the area where overlap seems to be maintained is in the area above the head. In an experiment, rats were placed in an enclosed area with a "shelter", where they could retreat when threatened. It was found that showing a threat, like a eyes. The result is that the rat threatened. It was found that showing a threat, like a shadow, from the sides of the enclosure did not result in response, but showing the threat from above led to the rats scampering into the shelter.

rats scampeting into un-shelter. Despite the uncoordinated eye movements, rodents do have the capacity for stereoscopic vision and display impressive object recognition. The mechanism of combining the overlap in images may be momentary orientation of the head to match the images or a computational method to make use of the two images. computational method to make use of the two images

But the overall object of how the rodent control of the now the rodent control of the eyes has evolved seems to be not to use both eyes to see visual targets while moving, but to have binocular vision only in the region overhead, which is where danger is most likely to lurk.

The writer can be contacted at simplescience@gmail.com









Making India strong and self-reliant with Technology and Innovation, equipping India's Armed Forces with state of the art weapons, systems and technology solutions.

DRDO's technologies have placed India among world's elite groups of:

- > 4 nations to have multi-level Strategic Deterrence capabilities
- 5 nations to have its own Ballistic Missile Defence (BMD) program
- 6 nations to have developed a Nuclear -Powered Submarine
- 7 nations to have developed own Main Battle Tank and 4th generation Combat Aircraft

DRDO continues in its march over path of excellence in developing cutting edge defence technologies.



Ministry of Defence, Govt. of India www.drdo.gov.in

Issued by Directorate of Public Interface

davp 10301/13/0178/1314



